

Equine Health Solutions

Keith L. Wagner DVM

18011 E State Route 58 Raymore, MO 64083 816-322-8040

www.equinehealthsolutions.com

EHS Recommended Foal Vaccination Schedule

<p>Foals born to mares vaccinated 4-6 weeks prior to foaling.</p> <ul style="list-style-type: none">• <u>6 months</u><ul style="list-style-type: none">○ Sleeping Sickness (1)○ Tetanus (1)○ Rabies (1) No earlier than 6 mo of age○ West Nile (1)• <u>7 months</u><ul style="list-style-type: none">○ Sleeping Sickness (2)○ Tetanus (2)○ Rabies (2)○ West Nile (2)• <u>8 months</u><ul style="list-style-type: none">○ Sleeping Sickness (3)○ Tetanus (3)• <u>9 months</u><ul style="list-style-type: none">○ Equine Infuenza (1)○ Equine Rhinopneumonitis (1)• <u>10 months</u><ul style="list-style-type: none">○ Equine Infuenza (2)○ Equine Rhinopneumonitis (2)• <u>11 months</u><ul style="list-style-type: none">○ Equine Infuenza (3)○ Equine Rhinopneumonitis (3)• <u>10-12 months</u><ul style="list-style-type: none">○ Rabies (3)○ West Nile (3) prior to mosquito season <p>Annual and semi-annual vaccinations there after.</p>	<p>Foals born to unvaccinated or unknown history mares.</p> <ul style="list-style-type: none">• <u>3 months</u><ul style="list-style-type: none">○ Sleeping Sickness (1)○ Tetanus (1)○ Rabies (1)○ West Nile (1)• <u>4 months</u><ul style="list-style-type: none">○ Sleeping Sickness (2)○ Tetanus (2)○ Rabies (2)○ West Nile (2)• <u>5 months</u><ul style="list-style-type: none">○ Sleeping Sickness (3)○ Tetanus (3)○ West Nile (3)• <u>6 months</u><ul style="list-style-type: none">○ Equine Infuenza (1)○ Equine Rhinopneumonitis (1)• <u>7 months</u><ul style="list-style-type: none">○ Equine Infuenza (2)○ Equine Rhinopneumonitis (2)• <u>8 months</u><ul style="list-style-type: none">○ Equine Infuenza (3)○ Equine Rhinopneumonitis (3)• <u>10-12 months</u><ul style="list-style-type: none">○ Rabies (3) <p>Annual and semi-annual vaccinations there after.</p>
--	---

Proper Foal De-Worming Schedule

- A foal should be de-wormed within 30 days of life and then every 60 days until 6 months of age.
- Panacur (fenbendazole) is recommended for foals until 6 months of age. When giving a dose, give double the weight of the foal.
- After 6 months of age the foal is ready to start adult de-wormer. Due to the development of resistant intestinal parasite population, it is recommended to de-worm the horse according to fecal test results and environmental evaluation.
- Foals that are 4 months of age that have **NOT** been de-wormed should consult a veterinarian to receive special de-worming instructions.